

The Construct Validity of the Illness Cognition Questionnaire: The Robustness of the Three-factor Structure Across Patients with Chronic Pain and Chronic Fatigue

Emelien Lauwerier · Geert Crombez ·
Stefaan Van Damme · Liesbet Goubert ·
Dirk Vogelaers · Andrea W. M. Evers

Published online: 15 September 2009
© International Society of Behavioral Medicine 2009

Abstract

Background The Illness Cognition Questionnaire (ICQ; Evers et al., *J Consult Clin Psychol*, 69:1026-1036, 2001) assesses three ways of cognitively evaluating the stressful and aversive character of a chronic illness: helplessness, acceptance, and perceived benefits.

Purpose The purpose of this study was to evaluate the construct validity of the ICQ in individuals with chronic pain and patients with chronic fatigue.

Method The ICQ was administered to 821 individuals with chronic pain and 295 patients with chronic fatigue. Confirmatory factor analyses were performed to assess the hypothesized three-factor structure, containing the factors “helplessness,” “acceptance,” and “perceived benefits.” A multigroup analysis was performed to investigate the stability of the factor structure in both groups.

Results Results confirmed the three-factor structure in the two samples. The factor structure was invariant across individuals with chronic pain and chronic fatigue.

Conclusion As the three-factor structure provided a good fit in both groups, we confirm the usefulness of the subscale scores in research and clinical practice.

Keywords Chronic pain · Chronic fatigue · Factor structure · Illness Cognition Questionnaire (ICQ)

Introduction

Medically unexplained symptoms, including many forms of chronic pain and chronic fatigue, often have a negative impact on quality of life, affecting physical, psychological, cognitive, and social domains of functioning [1, 2]. Symptom severity is often insufficient to fully explain their adverse effects upon functioning. Many variables have been identified that may hinder or promote adjustment to a life with chronic symptoms, such as self-efficacy, coping strategies, and illness beliefs [3, 4]. Although there is a wide variety of illness beliefs and cognitions [5–7] that are relevant for the adjustment to chronic symptoms, in this paper, we focus upon the three generic constructs of helplessness, acceptance, and experienced benefits of illness.

Some studies found evidence for the beneficial effects of perceived control over symptoms [8, 9]. Conversely, the repeated experience of lack of control over aversive events may result in helplessness [10]. In line with this, several studies have found that a perceived lack of control is associated with unfavorable outcomes in chronic pain conditions [11, 12].

Although there is merit in the idea that having control over symptoms may be associated with better adjustment, research in this respect has yielded inconclusive results [13]. As an example, it has been found that attempting to control or solve pain when actual control is low may increase fear, worry, catastrophic thinking, and hypervigilance [14, 15]. In some situations, abandoning the struggle to control symptoms and accepting the illness may be more

E. Lauwerier · G. Crombez (✉) · S. Van Damme · L. Goubert
Department of Experimental–Clinical and Health Psychology,
Ghent University,
Henri Dunantlaan 2,
9000 Ghent, Belgium
e-mail: Geert.Crombez@UGent.be

D. Vogelaers
General Internal Medicine and Infectious Diseases,
Ghent University Hospital,
Ghent, Belgium

A. W. M. Evers
Department of Medical Psychology, University of Nijmegen,
Nijmegen, the Netherlands

adaptive. This idea has been much less studied than the role of perceived control. Nonetheless, acceptance is part of several coping models [16–18]. It has been found that patients who are accepting of pain reported less depression, anxiety, and disability [19]. In a questionnaire study in chronic pain patients, Viane et al [20] showed that acceptance was related to better psychological, but not physical, well-being. In agreement with the above findings, research in patients with chronic fatigue has pointed out that acceptance was related to more emotional stability and less psychological distress [21].

To reduce the emotional strain of adversity, one can also try to make sense out of the losses the situation has caused. Rather than downgrading the importance of the blocked goal and accepting the constraints of the situation, one can give a new, positive evaluation to the aversive situation. Affleck and Tennen [22] were one of the first to underline the importance of “benefit finding” or seeing positive side effects in otherwise aversive situations. It has already been found that these illness benefits buffer negative effects of perceived health stresses on subjective well-being [23].

Several self-report measures have been developed to assess the above-reported concepts of helplessness, acceptance, and benefit finding. Some of these instruments are symptom specific [19, 24, 25] and, hence, do not allow comparisons between groups with different symptoms. An example of a disease-specific instrument is the Perceived Control and Benefits Questionnaire [25], designed to measure personal control and the perception of benefits in patients with chronic pain. Other measures assess the constructs in a trait-like fashion unrelated to specific situations such as chronic illnesses [26–28]. For example, the Coping with Problems Experienced inventory [27] assesses coping strategies as relatively stable preferences. With this type of measures, situational influences or constraints may be easily overlooked.

Recently, the Illness Cognition Questionnaire (ICQ) has been developed as a generic measure of illness beliefs [29]. The instrument consists of three subscales, i.e., helplessness, acceptance, and perceived benefits. It allows comparison across chronic conditions. Furthermore, items and instructions are not formulated in a trait-like manner, potentially allowing us to assess situational influences. Evers et al [29] investigated the psychometric properties (i.e., reliability and validity) of the ICQ. They conducted a principal components analysis with oblique rotation on ICQ scores derived from 263 patients with rheumatoid arthritis and obtained a three-factor solution accounting for 62% of the total variance. The factors were labeled helplessness, acceptance, and perceived benefits. They also performed a confirmatory factor analysis on ICQ scores obtained from 167 patients with multiple sclerosis. Analyses indicated that the three-factor structure of the ICQ provided a satisfactory

fit to the data. In sum, the ICQ showed a strong internal consistency, reliability, and good construct and predictive validity. Helplessness was associated with unfavorable changes, whereas acceptance and perceived benefits were related to beneficial changes in physical and psychological health on the long term.

Although the ICQ has shown good psychometric qualities, the question remains whether the factor structure can be replicated in patient groups with medically unexplained symptoms, such as chronic pain and chronic fatigue. Therefore, the aims of this study were twofold: First, we investigated the construct validity of the Illness Cognition Questionnaire by means of a confirmatory factor analysis in samples of individuals with chronic pain and chronic fatigue. Second, we investigated the stability of the factor structure across the two groups using a multisample analysis. To invigorate further research and clinical practice, we provided norms for the subscales of the ICQ.

Method

Participants

Data were collected from two samples which completed the same Dutch version of the ICQ. For both samples, no data are available on response rate and reasons for nonparticipation. Additionally, participants received no reward for participation.

The first sample consisted of 871 Dutch individuals with chronic pain who were a member of one of two self-help groups. For fifty patients, some item scores were missing, reducing our sample to 821 patients (19% males; 81% females), aged between 19 and 99 years (mean age=50.75 years, SD=10.4). The average pain duration was 183 months (SD=131, range 12–732). Most patients reported pain at multiple sites (45.9%) or back pain (28.8%). The majority of the entire sample reported secondary education as highest education level (66.9%), whereas only small groups received either a lower education (6.5%) or a higher education (longer than the age of 18; 26.6%). Within the entire sample, the majority was married or living together (76.3%).

After reduction because of incomplete data, the second sample consisted of 295 Flemish chronic fatigue patients, who were on a waiting list for cognitive behavioral therapy and fulfilled all the “Center for Disease Control and Prevention” criteria for chronic fatigue syndrome [30]. The sample consisted of 12.5% men and 87.5% women, aged between 18 and 64 (mean age=40.32 years, SD=8). A small group reported a lower education as highest education level (8.3%); 56.3% had secondary education, and 35.4% received a higher education.

Measures

Illness Cognitions

The ICQ [29] was used to measure helplessness, acceptance, and perceived benefits. This is a 18-item questionnaire that contains three six-item scales related to the factors helplessness, acceptance, and perceived benefits, each with a scoring range of 6–24 (e.g., helplessness: “My illness limits me in everything that is important to me”, “My illness frequently makes me feel helpless”; acceptance: “I have learned to live with my illness”, “I can accept my illness well”; perceived benefits: “Dealing with my illness has made me a stronger person”, “My illness has taught me to enjoy the moment more”). Each item is answered on a four-point Likert scale to the extent to which one agrees with the item (1 = *not at all*, 2 = *somewhat*, 3 = *to a large extent*, 4 = *completely*).

Exploratory factor analysis in a sample of patients with rheumatoid arthritis ($n=263$) revealed a three-factor solution accounting for 62% of the variance. Confirmatory factor analysis in a sample of patients with multiple sclerosis ($n=167$) has confirmed the assumed three-factor structure. Cronbach’s alpha demonstrated adequate internal consistencies for all scales, ranging from 0.84 to 0.91 in both samples. Pearson’s correlation coefficients between two administrations of the questionnaire with a 1-year time interval were all above 0.67, indicating good test–retest reliability for all scales in both samples. Evidence has also been found for good concurrent and predictive validity [29].

Analytical Strategy

Confirmatory factor analyses were performed using AMOS 7.0 [31]. The hypothesized three-factor structure was tested in the chronic pain sample. The chronic fatigue sample was used to cross-validate the structure. The fit of the model was estimated with the maximum likelihood algorithm while allowing the latent variables to correlate. Each item was assumed to load only on one factor. Subsequently, a multigroup analysis was carried out in order to examine the invariance of the factor structure across both groups. Model fit is usually evaluated using the χ^2 goodness-of-fit statistic and several fit indices [32]. As recommended by Hu and Bentler [33], we used a strategy combining the following fit indices: the standardized version of Jöreskog and Sörbom’s [34] root mean square residual (SRMR) [35] and the comparative fit index (CFI) [36].

The χ^2 goodness-of-fit statistic assesses the overall fit of the model and, in particular, whether a significant amount of observed covariance between items remains unexplained by the model. A significant χ^2 is indicative of a bad model fit. The main shortcoming of this fit index is its sensitivity

to sample size. In a small sample, a poor fit may result in a χ^2 that is nonsignificant. Equally, it is also possible that, in large samples, a good fit results in a statistically significant χ^2 [37]. Therefore, for further interpretation, we rather looked at the fit index χ^2/df (CMIN/DF), which is the minimum sample discrepancy divided by the degrees of freedom. According to Marsh and Hovecar [38], CMIN/DF values between 2 and 5 represent a reasonable model fit.

The standardized version of the Jöreskog and Sörbom’s [34] SRMR [35] is an absolute fit index, assessing how well an a priori model reproduces the sample data. Hu and Bentler [33] found that the SRMR is the most sensitive fit index for models with misspecified factor covariances. Values close to 0 suggest that the data fit the model.

To measure the proportionate improvement in model fit by comparing the target model with a baseline model, we used the CFI. The CFI is one of the most sensitive indices to models with misspecified factor loading(s). The CFI usually ranges between 0 and 1, with values above 0.95 indicating a good fit [33].

We used a combination of indices to evaluate model fit, in particular CFI and SRMR. A model has a good fit when the CFI value is close to 0.95 or larger and when the SRMR value is close to 0.09 or lower. In line with Hu and Bentler [33], we used the following criteria: for a good model fit, CFI>0.94 and SRMR<0.09; for an adequate model fit, CFI>0.90 and SRMR<0.09; and for a poor model fit, CFI<0.90 and SRMR>0.09.

Results

Descriptive and Correlational Statistics

Table 1 reports the means, standard deviations, and internal consistencies of the subscales and Pearson correlation coefficients between the subscales of the ICQ. The internal consistencies of all subscales in the two groups were good (range=0.81–0.91). Overall, the pattern of correlations among the subscales of helplessness, acceptance, and perceived benefits was as expected. Helplessness was negatively related to the two other constructs acceptance and perceived benefits, respectively. However, the association with perceived benefits was less pronounced. In contrast, acceptance was positively related to perceived benefits. There were no significant age effects, except for a positive correlation between helplessness and age in chronic pain patients ($r=0.16$, $p<0.001$), indicating greater helplessness among older patients in the pain sample.

Confirmatory Factor Analyses

The standardized factor loadings of the three-factor model for the chronic pain and chronic fatigue sample are

Table 1 Means (*M*), standard deviations (*SD*), internal consistency (α), Pearson correlation coefficients, and their 95% confidence intervals among ICQ subscales: helplessness, acceptance, and perceived benefits

	<i>M</i> (<i>SD</i>)	α	2	3
Chronic pain				
Helplessness (ICQ)	14.52 (4.30)	0.88	-0.48*** [-0.53; -0.43]	-0.07 [-0.014; 0]
Acceptance (ICQ)	14.60 (4.13)	0.91	–	0.44*** [0.38; 0.50]
Perceived benefits (ICQ)	15.02 (4.31)	0.83	–	–
Chronic fatigue				
Helplessness (ICQ)	16.57 (3.86)	0.83	-0.53*** [-0.61; -0.45]	-0.29*** [-0.39; -0.18]
Acceptance (ICQ)	11.95 (3.73)	0.90	–	0.48*** [0.39; 0.57]
Perceived benefits (ICQ)	12.50 (4.09)	0.81	–	–

ICQ Illness Cognition Questionnaire

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

presented in Fig. 1. As for the individuals with chronic pain, although results showed a significant χ^2 (χ^2 (*df*)=683.176 (132), $p < 0.001$, CMIN/DF=5.20), the fit indices supported the adequate fit of the model (CFI=0.93, SRMR=0.06). In the chronic fatigue sample, we found significant overall fit (χ^2 (*df*)=326.84 (132), $p < .001$, CMIN/DF=2.48) of the three-factor model to the data. The fit indices indicated an adequate fit (CFI=0.92, SRMR=0.07). After inspection of the modification indices, a correlated residual between items 7 (“My illness makes me feel useless at times”) and 15 (“My illness frequently makes me feel helpless”) was detected in both samples. This finding indicates that these items have something in common, which is not reflected by the remaining items that load on the helplessness subscale. The residual may probably be due to content overlap between the items. In both samples, the model was refitted to the data, allowing a free estimation of the error covariance between items 7 and 15. The improvement in fit provided by the model with residual correlation between items 7 and 15 was significant in individuals with chronic pain ($\Delta\chi_{(1)}^2=71.37$, $p < 0.001$, CMIN/DF=4.70) and in individuals with chronic fatigue ($\Delta\chi_{(1)}^2=27.90$, $p < 0.001$, CMIN/DF=2.30). After refitting the model, the three-factor structure had an adequate fit to the data, both in the chronic pain sample (CFI=0.94, SRMR=0.06) and in the chronic fatigue sample (CFI=0.93, SRMR=0.07). Inspection of the modification indices indicated that other minor improvements were possible in both samples. It was decided not to include these changes because of model parsimony.

Invariance of the Factor Structure

To examine whether the three-factor structure is invariant across the two chronic conditions, a multigroup analysis was performed. A restrictive model [39], equating the number of factors, the factor loadings, the correlations




Fig. 1 Standardized factor loadings as obtained with confirmatory factor analysis shown for chronic pain patients and patients with chronic fatigue (between parentheses)

between the factors and the error variances was investigated. The overall fit was significant (χ^2 (df)=1,143.800 (303), $p < 0.001$, CMIN/DF=3.78). The fit statistics for the restrictive model reflected an adequate fit to the data (CFI=0.93, SRMR=0.06).

Norms

Levene's test for equality of variances was conducted in order to examine if equal variances could be assumed. Because equal variances were assumed for the subscale perceived benefits, we further used the t test for equality of variances in order to analyze differences in scores. For the subscales helplessness and perceived benefits, equality of variances could not be assumed. Therefore, we used a t test for inequality of variances to analyze differences in scores in those subscales. Analyses revealed significant differences for the scores on helplessness ($t(574.5) = -7.59$, $p < 0.001$), acceptance ($t(570.25) = 10.16$, $p < 0.001$), and perceived benefits ($t(1,114) = 8.70$, $p < .001$) between individuals with chronic pain ($n = 821$) and individuals with chronic fatigue ($n = 295$). As a result, separate norms were calculated for the two different chronic conditions (see Table 2).

When examining gender differences, we found a significant difference in scores on acceptance ($t(731) = 2.68$, $p < 0.01$), with men scoring higher ($M = 14$, $SD = 4$) than women ($M = 13$, $SD = 4$). Scores on helplessness ($t(731) = 0.144$, ns) and perceived benefits ($t(272.42) = -0.64$, ns) did not differ between the two sexes. To investigate whether the gender effect for acceptance was due to the type of medically unexplained complaint (chronic pain versus

chronic fatigue), we performed an (group \times gender) analysis of variance (ANOVA) upon the subscale acceptance. The ANOVA showed a significant effect of group, $F(1, 729) = 25.032$, $p < 0.001$. The main effect of gender was not significant, $F(1, 729) = 1.113$, $p = 0.29$. There was also no interaction effect between condition and gender, $F(1, 729) = 0.087$, $p = 0.77$. Results seem to indicate that there is no effect of gender on acceptance.

Discussion

The present study investigated the construct validity of the ICQ [29] using confirmatory factor analysis in two different samples, namely individuals with chronic pain and individuals with chronic fatigue. Furthermore, the invariance of the factor structure across the two groups was examined using a multigroup analysis.

The goodness-of-fit indices of the three-factor solution indicated an adequate fit to the data in both samples. Furthermore, although the chronic pain and chronic fatigue samples differed with regard to the mean scores on the subscales, the factor structure proved to be invariant across the two samples. It seems likely to assume that equivalent illness beliefs, reflected by the subscales of the ICQ, are tapped in both conditions. To our knowledge, the current study is the first to support the invariance of the factor structure of the ICQ across different samples with medically unexplained symptoms.

Concerning the psychometric properties of the Illness Cognition Questionnaire, results demonstrated adequate internal consistencies for all scales. Intercorrelations be-

Table 2 Deciles and quartiles for the chronic pain ($n = 821$) and chronic fatigue sample ($n = 295$)

	Chronic pain			Chronic fatigue		
	Helplessness	Acceptance	Perceived benefits	Helplessness	Acceptance	Perceived benefits
Deciles						
10	9	9	9	7	11.6	7
20	11	11	11	8	13	9
30	12	12	12	10	14	10
40	13	13	14	11	16	11
50	14	14	15	12	17	12
60	15	16	16	12	18	13
70	17	17	17	13.2	19	15
80	18	18	19	15	20	16
90	21	20	21	17	22	18
Quartiles						
25	11	12	12	9	13	9
50	14	14	15	12	17	12
75	18	18	18	14	19	15

tween the scales were in line with the expectations and at the same time insignificant to moderate, which revealed their content validity. Based on these findings, we confirm the usefulness of the subscale scores in research and clinical practice. Our norms may be used to describe patient samples in terms of the illness cognitions. Differences in cognitions can be expected between samples from different settings. Furthermore, clinicians may use the upper quartile cutoff scores of the illness cognitions for screening and diagnostic purposes.

The results of this study have a number of implications. First, they give support to the idea that the ICQ is an instrument that can be used for individuals with different medically unexplained symptoms. Second, although there are large differences in nature and experience of complaints between chronic pain and chronic fatigue, the ICQ seems to assess the same processes of illness beliefs. Thus, differences in subscale scores between those samples are rather quantitative than qualitative in nature. Third, our results are in line with the idea that acceptance and perceived benefits play a role in the adaptation to uncontrollable symptoms. We endorse the view of Rothermund [40] stating that, although somewhat neglected in the past, we cannot overlook the importance of disengagement and acceptance in situations where control is low or nonexistent.

The strength of the current study lies in the large sample size for both the chronic pain and chronic fatigue group. That way, the study has adequate statistical power and results in more accurate estimates of the statistics. There are a number of limitations to this study. First, more studies are needed in individuals with chronic pain and chronic fatigue to investigate whether the results generalize to other samples and to investigate the relationships between the ICQ scales and health-related outcomes. Second, prospective studies are needed to examine the predictive validity of the ICQ scales in long-term adaptation to chronic illness. Third, the stability over time (i.e., test–retest reliability) was not investigated in the present study and should be looked upon in future studies. Fourth, further research may examine if certain illness beliefs are involved in the maintenance of chronic suffering. There is evidence that, in chronic pain patients, repeating attempts to control or solve the pain problem may fuel negative consequences [14]. Beliefs related to the loss of control, like helplessness, may be more prominent in patients who continuously fail in finding a solution for their symptoms, like pain or fatigue. Conversely, the acceptance of illness may buffer against the adverse impact of symptom severity. Finally, research is needed to investigate to what extent treatment is able to change these cognitions and to what extent these changes mediate treatment success.

Acknowledgements We want to thank the Belgian Pain League who coordinates several self-help groups for pain patients in Belgium, the National Association for patients with Fibromyalgia of the Netherlands (FES; Nationale Vereniging voor Fibromyalgie-patiënten “Eendrachtig-Sterk”), and the psychosomatic center of University Hospital Ghent for collection of the data.

References

1. Anderson JS, Ferrans CE. The quality of life of persons with chronic fatigue syndrome. *J Nerv Ment Dis.* 1997;185:359–67.
2. Niv D, Kreitler S. Pain and quality of life. *Pain Pract.* 2001;1:150–61.
3. Keefe FJ, Rumble ME, Scipio CD, Giordano LA, Perri LM. Psychological aspects of persistent pain: current state of the science. *J Pain.* 2004;5:195–211.
4. Vercoulen JHMM, Swanink CM, Galama JM, Fennis JF, Jongen PJ, Hommes OR, et al. The persistent of fatigue in chronic fatigue syndrome and multiple sclerosis: development of a model. *J Psychosom Res.* 1998;45:507–17.
5. Devellis BM, Blalock SJ. Illness attributions and hopelessness depression: the role of hopelessness expectancy. *J Abnorm Psychol.* 1992;101:257–64.
6. Leventhal H, Brissette I, Leventhal EA. The common-sense model of self-regulation of health and illness. In: Cameron LD, Leventhal H, editors. *The self-regulation of health and illness behaviour.* London: Routledge; 2003. p. 42–65.
7. Weinman J, Petrie KJ, Moss-Morris R, Horne R. The illness perception questionnaire: a new method for assessing the cognitive representation of illness. *Psychol Health.* 1996;11:431–45.
8. Buckelew SP, Parker JC, Keefe FJ, Deuser WE, Crews TM, Conway R, et al. Self-efficacy and pain behavior among subjects with fibromyalgia. *Pain.* 1994;59:377–84.
9. Jensen MP, Karoly P. Control beliefs, coping efforts, and adjustment to chronic pain. *J Consult Clin Psychol.* 1991;59: 431–8.
10. Overmier JB, Seligman MEP. Effects of inescapable shock upon subsequent escape and avoidance learning. *J Comp Physiol Psychol.* 1967;63:23–33.
11. Koleck M, Mazaux JM, Rasclé N, Bruchon-Schweitzer M. Psycho-social factors and coping strategies as predictors of chronic evolution and quality of life in patients with low back pain: a prospective study. *Eur J Pain.* 2006;10:1–11.
12. Nicassio PM, Schuman C, Radojevic V, Weisman MH. Helplessness as a mediator of health status in fibromyalgia. *Cognit Ther Res.* 1999;23:181–96.
13. McCracken LM, Eccleston C. Coping or acceptance: what to do about chronic pain? *Pain.* 2003;105:197–204.
14. Crombez G, Eccleston C, De Vlieger P, Van Damme S, De Clercq A. Is it better to have controlled and lost than never to have controlled at all? An experimental investigation of control over pain. *Pain.* 2008;137:631–9.
15. Eccleston C, Crombez G. Worry and chronic pain: a misdirected problem solving model. *Pain.* 2007;132:233–6.
16. Brandstädter J, Renner G. Tenacious goal pursuit and flexible goal adjustment: explication and age-related analysis of assimilative and accommodative strategies of coping. *Psychol Aging.* 1990; 5:58–97.
17. Heckhausen J, Schulz R. A life-span theory of control. *Psychol Rev.* 1995;102:284–304.
18. Scheier MF, Carver CS. Goals and confidence as self-regulatory elements underlying health and illness behavior. In: Cameron LD,

- Leventhal H, editors. *The self-regulation of health and illness behaviour*. London: Routledge; 2003. p. 17–41.
19. McCracken LM. Learning to live with the pain: acceptance of pain predicts adjustment in persons with chronic pain. *Pain*. 1998;74:21–7.
 20. Viane I, Crombez G, Eccleston C, Poppe C, Devulder J, Van Houdenhove B, et al. Acceptance of pain is an independent predictor of mental well-being in patients with chronic pain: empirical evidence and reappraisal. *Pain*. 2003;106:65–72.
 21. Van Damme S, Crombez G, Van Houdenhove B, Mariman A, Michielsens W. Well-being in patients with chronic fatigue syndrome: the role of acceptance. *J Psychosom Res*. 2006;61:595–9.
 22. Affleck T, Tennen H. Construing benefits from adversity: adaptational significance and dispositional underpinnings. *J Pers*. 1996;64:899–922.
 23. Wrosch C, Heckhausen J, Lachman ME. Primary and secondary control strategies for managing health and financial stress across adulthood. *Psychol Aging*. 2000;15:387–99.
 24. Flor H, Behle DJ, Birbaumer N. Assessment of pain-related cognitions in chronic pain patients. *Behav Res Ther*. 1993;31:63–73.
 25. Tennen H, Affleck G, Urrows S, Higgins P, Mendola R. Perceiving control, construing benefits, and daily processes in rheumatoid arthritis. *Can J Behav Sci*. 1992;24:186–203.
 26. Beck AT, Weissman A, Lester D, Trexler L. The measurement of pessimism: the hopelessness scale. *J Consult Clin Psychol*. 1974;42:861–5.
 27. Carver CS, Scheier MF, Weintraub JK. Assessing coping strategies: a theoretically based approach. *J Pers Soc Psychol*. 1989;56:267–83.
 28. Millon T, Green CJ, Meagher RB. A new psychodiagnostic tool for clients in rehabilitation settings: the MBHI. *Rehabil Psychol*. 1982;27:23–35.
 29. Evers AWM, Kraaimat F, van Lankveld W, Jongen PJH, Jacobs JWG, Bijlsma JWJ. Beyond unfavorable thinking: the illness cognition questionnaire for chronic disease. *J Consult Clin Psychol*. 2001;69:1026–36.
 30. Fukuda K, Straus SE, Hickie I, Sharpe MC, Dobbins JG, Komaroff A. The chronic fatigue syndrome: a comprehensive approach to its definition and study. *Ann Intern Med*. 1994;121:953–9.
 31. Arbuckle JL. Amos (version 7.0) [Computer Program]. Chicago: SPSS; 2006.
 32. Bollen KA, Long JS. Introduction. In: Bollen KA, Long JS, editors. *Testing structural equation models*. Newbury Park: Sage; 1993. p. 1–9.
 33. Hu LT, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Struct Equ Modeling*. 1999;6:1–55.
 34. Jöreskog KG, Sörbom D, Lisrel V. *Analysis of linear structural relationship by the method of maximum likelihood*. Chicago: National Educational Resources; 1981.
 35. Bentler PM (1995) *EQS Structural equations program manual*. Encino, CA: Multivariate Software
 36. Bentler PM. Comparative fit indexes in structural models. *Psychol Bull*. 1990;107:238–46.
 37. Marsh HW, Balla JR, McDonald RP. Goodness-of-fit indexes in confirmatory factor analysis: the effect of sample size. *Psychol Bull*. 1988;103:391–410.
 38. Marsh HW, Hovecar D. Application of confirmatory factor analysis to the study of self-concept: first- and higher order factor models and their invariance across groups. *Psychol Bull*. 1985;97:562–82.
 39. Hoyle RH, Smith GT. Formulating clinical research hypotheses as structural equation models: a conceptual overview. *J Consult Clin Psychol*. 1994;62:429–40.
 40. Rothermund K. Hanging on and letting go in the pursuit of health goals: psychological mechanisms to cope with a regulatory dilemma. In: De Ridder D, De Wit J, editors. *Self regulation in health behaviour*. Chichester (UK): Wiley; 2006. p. 217–41.